

2X Hi G9 Taq PCR Master Mix

#G4804 100 Reactions

Store at -20°C

Spin tubes briefly before use

Catalogue Number	Pack Size	☑
G4804	100 Reaction	✓
G4804A	250 Reaction	
G4804B	1000 Reaction	

Introduction

Codon optimized Hi G9 Taq DNA Polymerase gene of *Thermus aquaticus* was cloned and purified from *E. coli* host. The enzyme consists of a single polypeptide with a molecular weight of approximately 94 kDa. Hi G9 Taq DNA polymerase is heat-stable and will synthesize DNA at elevated temperatures from single-stranded templates in the presence of primer.

Guidelines for using Prime Taq DNA Polymerase

The Master Mix contains Hi-G9 Taq DNA polymerase, Hi G9 Taq DNA reaction buffer, 5mM MgCl₂, and 0.4mM of each dNTP.

Package information

G4804 (100 Reactions)	Material provided: 2X Hi G9 Taq PCR Master Mix (2 x 1.25 ml), 10ul Control DNA template, 10ul Control primer Mix
G4804A (250 Reactions)	Material provided: 2X Hi G9 Taq PCR Master Mix (5 x 1.25ml), 10ul Control DNA template, 10ul Control primer Mix
G4804B (1000 Reactions)	Material provided: 2X Hi G9 Taq PCR Master Mix (20 x 1.25ml), 10ul Control DNA template, 10ul Control primer Mix

Unit definition

One unit incorporates 10nmol of deoxy ribonucleotide into acid-insoluble product in 30 minutes at 74°C. Unit assay conditions: 25 mM TAPS (pH 9.3), 50 mM KCl, 2 mM MgCl₂, 1 mM DTT, 0.2 mM dATP, dCTP, dGTP, dTTP utilizing M13mp18DNA as template.

Enzyme

The amount of enzyme will be 1 Unit in 50 µl reactions finally. When cloning fragments amplified with Hi G9 Taq DNA Polymerase an extra adenine overhang base is incorporated at the 3' end of the amplified PCR product. That A overhang is required to perform the TA cloning.

Table 1. Pipetting instructions

(add items in this order).

Component	50 µl Reaction	Final conc.
H ₂ O	To 50µl	
2x PCR Master Mix	25 µl	1X
dNTPs	-	0.2mM each
MgCl ₂	-	2.5mM
Forward primer	X µl	0.5µM
Reverse Primer	X µl	0.5µM
Template DNA	X µl	

In some template to get optimum result primers conc and MgCl₂ conc. needed to be optimized

Table 2. Cycling Instructions

Cycle step	2-step Protocol		Cycles
	Temp.	Time	
Initial Denaturation	94°C	5min	1
Denaturation	94°C	15-30sec	25-35
Annealing	T _m	15-60sec	
Extension	X°C (see notes)	60sec/1kb	
Final extension	X°C	10 min	1
	4°C	Hold	

Notes:

- Use 94°C for denaturation.
- Use 15–60 s/kb for extension..
- The annealing temp depends on the primer.
- Use extension at 72°C for upto 6kb amplification and at 68°C for above 6kb amplification.
- Note: Hi G9 Taq DNA Polymerase produces A overhang at the 3' end.

Template

For low complexity DNA (e.g. plasmid, lambda or BAC DNA): 2 pg–20 ng per 50 µl reaction volume needed to be used. For high complexity genomic DNA, the amount of DNA template should be 20–200 ng per 50 µl reaction volume. If cDNA synthesis reaction mixture is used directly as a source for the template, the volume used should not exceed 10 % of the final PCR reaction volume.

Primers

The recommendation for final primer concentration is 0.5 µM. If required, the primer concentration may be optimized between 0.2–1.0 µM. The results from primer T_m calculations can vary significantly depending on the method used. Always use the T_m calculator and instructions from reputed website to determine the T_m

values of primers and optimal annealing temperature. If using a two-step PCR protocol, where both primer annealing and extension occur in a single step at 72°C, the primers should be designed accordingly.

Mg²⁺ and dNTP

2X Hi G9 Taq PCR Mater Mix is optimized to work well for most amplicons. Standard concentration of MgCl₂ in final reaction condition is 2.5 mM.

DMSO

The recommended reaction conditions for GC-rich templates include 3 % DMSO as a PCR additive, which aids in the denaturing of templates with high GC contents. For further optimization the amount of DMSO should be increased in 2 % increments. In some cases DMSO may also be required for supercoiled plasmids to relax for denaturation. Other PCR additives such as formamide, glycerol, and betaine are also compatible with Hi G9 Taq DNA Polymerase. If high DMSO concentration is used, the annealing temperature must be decreased, as DMSO affects the melting point of the primers. It has been reported that 10 % DMSO decreases annealing temperature by 5.5–6.0°C.

Troubleshooting

Optimize annealing temperature.	Increase extension time.
<ul style="list-style-type: none"> Repeat and make sure that there are no pipetting errors. 	<ul style="list-style-type: none"> Increase the number of cycles.
<ul style="list-style-type: none"> Make sure the cycling protocol was performed as recommended. 	<ul style="list-style-type: none"> Titrate DMSO (2–8 %) in the reaction.
<ul style="list-style-type: none"> Use fresh high-quality dNTPs. 	<ul style="list-style-type: none"> Denaturation temperature may be too low. Optimal denaturation temperature for most templates is 94–98°C.
<ul style="list-style-type: none"> Do not use dNTP mix or primers that contain dUTP or dITP. 	<ul style="list-style-type: none"> Optimize denaturation time.
<ul style="list-style-type: none"> Titrate template amount. Template DNA may be damaged. Use freshly prepared template. 	<ul style="list-style-type: none"> Check the purity and concentration of the primers. Check primer design.
Non-specific products - High molecular weight smears	
<ul style="list-style-type: none"> Make sure the extension time used was not too long. 	<ul style="list-style-type: none"> Titrate template amount.
<ul style="list-style-type: none"> Reduce the total number of cycles. 	<ul style="list-style-type: none"> Optimize denaturation temperature.
<ul style="list-style-type: none"> Increase annealing temperature or try 2-step PCR protocol. 	<ul style="list-style-type: none"> Decrease primer concentration.

Non-specific products - Low molecular weight discrete bands

- | | |
|---|---|
| <ul style="list-style-type: none"> Increase annealing temperature | <ul style="list-style-type: none"> Titrate template amount. |
| <ul style="list-style-type: none"> Make sure the extension time used was not too long. | <ul style="list-style-type: none"> Decrease primer concentration. Design new primers. |

Component	Volume
2X Hi G9 Taq PCR Master Mix	25 µl
Control DNA template	1 µl
Control primer Mix	1 µl
Nuclease-free water	23 µl

Control PCR set-up

Cycling conditions	
Step 1	94°C – 5 min
Step 2	94°C – 30 s
Step 3	57°C – 30 s
Step 4	72°C – 4 min
Step 5	72°C – 10 min
Step 6	4 °C – Hold

Repeat Steps 2 to 4 for 40 Cycles. Control PCR will produce a 2Kb product.