

## G9 Taq DNA Polymerase 10X buffer with MgCl<sub>2</sub>

#G7115B 500 units

Store at -20°C

Catalogue Number	Pack Size	<input checked="" type="checkbox"/>
G7115	1000 Unit	
G7115B	500 Unit	<input checked="" type="checkbox"/>
G7115A	5000 Unit	

### Introduction

Codon optimized Taq DNA Polymerase gene of *Thermus aquaticus* was cloned and purified from *E. coli* host. The enzyme consists of a single polypeptide with a molecular weight of approximately 94 kDa. Taq DNA polymerase is heat-stable and will synthesize DNA at elevated temperatures from single-stranded templates in the presence of primer. It can be used for PCR of upto **6kb** product.

### Guidelines for using G9 Taq DNA Polymerase

G9 Taq DNA Polymerase is provided with **10x G9Taq DNA Reaction Buffer**. The buffer contains 2.5mM MgCl<sub>2</sub>. No separate tube of MgCl<sub>2</sub> is provided to maintain the flexibility.

### Package information

Component	G7115B	G7115	G7115A
Taq (2.5 unit/λ)	500 Unit	(2X 500 Unit)	(10X 500 Unit)
10X Taq Buffer with MgCl <sub>2</sub>	(2 X 1.25 ml)	(4X 1.25 ml)	(20X 1.25 ml)
Control DNA template	10 µl	10 µl	10 µl
Control Primer mix	10 µl	10 µl	10 µl

### Unit definition

One unit incorporates 10nmol of deoxy-ribonucleotide into acid-insoluble product in 30 minutes at 74°C. Unit assay conditions: 25 mM TAPS (pH 9.3), 50 mM KCl, 2 mM MgCl<sub>2</sub>, 1 mM DTT, 0.2 mM dATP, dCTP, dGTP, dTTP utilizing M13mp18DNA as template.

### Enzyme

The optimal amount of enzyme is 0.5 µl in 50 µl reactions. When cloning fragments amplified with G9 taq DNA Polymerase an extra adenine overhang base is incorporated at the 3' end of the amplified PCR product. That A overhang is required to perform the TA cloning.

**Table 1. Pipetting instructions** (add items in this order).

Component	50 reaction	Final conc.
Template DNA	X µl	<1,000 ng
10x Reaction Buffer with MgCl <sub>2</sub>	5µl	1X
10 mM dNTPs	1 µl	200µM each
Forward primer#	1 µl	0.2 µM
Reverse primer#	1 µl	0.2 µM
H <sub>2</sub> O	To 50 µl	
G9 Taq DNA Polymerase	0.5 µl	1.25U

### Protocol

- Gently vortex and briefly centrifuge all solutions after thawing.
- Place a thin-walled PCR tube on ice and add the components as mentioned in Table 1 for each 50 µL reaction.

**Table 2. Cycling Instructions**

Cycle step	2-step Protocol		Cycles
	Temp.	Time	
Initial Denaturation	94°C	5min	1
Denaturation	94°C	15-30sec	25-35
Annealing	T <sub>m</sub>	15-60sec	
Extension	68-72°C	2 min /1kb	
Final extension	68-72°C	5-15min	1

- Mix the samples and spin down as quickly as possible.
- Perform PCR using recommended thermal cycling conditions (see Table 2). For higher amplification (>4 Kb) extension at 68°C is preferable.

### Template

For low complexity DNA (e.g. plasmid, lambda or BAC DNA): 1 pg–10 ng per 25 µl reaction volume or 2 pg– 20 ng per 50 µl reaction volume. For high complexity genomic DNA, the amount of DNA template should be 10–400 ng per 25 µl reaction volume or 20–800 ng per 50 µl reaction volume.

If cDNA synthesis reaction mixture is used directly as a source for the template, the volume used should not exceed 10 % of the final PCR reaction volume.

### Mg<sup>2+</sup> and dNTP

G9 Taq reaction buffer is optimized to work well for most amplicons. Standard concentration of MgCl<sub>2</sub> in PCR reaction mixture is 2.5mM but to optimum concentration might vary between 1-5 mM. High quality dNTPs should be used for optimal performance with G9 Taq DNA Polymerase. The polymerase cannot read dUTP-derivatives or dITP in the template strand so the use of these analogues or primers containing them is not

recommended. For optimal results always use 200  $\mu\text{M}$  of each dNTP.

### DMSO

The recommended reaction conditions for GC-rich templates include 3 % DMSO as a PCR additive, which aids in the denaturing of templates with high GC contents. For further optimization the amount of DMSO should be increased in 2 % increments. In some cases DMSO may also be required for supercoiled plasmids to relax for denaturation. Other PCR additives such as formamide, glycerol, and betaine are also compatible with G9 Taq DNA Polymerase. If high DMSO concentration is used, the annealing temperature must be decreased, as DMSO affects the melting point of the primers. It has been reported that 10 % DMSO decreases annealing temperature by 5.5–6.0°C.

### Troubleshooting

Optimize annealing temperature.	Increase extension time.
<ul style="list-style-type: none"> <li>Optimize annealing temperature.</li> <li>Repeat and make sure that there are no pipetting errors.</li> </ul>	<ul style="list-style-type: none"> <li>Increase the number of cycles.</li> </ul>
<ul style="list-style-type: none"> <li>Make sure the cycling protocol was performed as recommended.</li> </ul>	<ul style="list-style-type: none"> <li>Titrate DMSO (2–8 %) in the reaction.</li> </ul>
<ul style="list-style-type: none"> <li>Use fresh high-quality dNTPs.</li> </ul>	<ul style="list-style-type: none"> <li>Denaturation temperature may be too low. Optimal denaturation temperature for most templates is 94–98°C.</li> </ul>
<ul style="list-style-type: none"> <li>Do not use dNTP mix or primers that contain dUTP or dITP.</li> </ul>	<ul style="list-style-type: none"> <li>Optimize denaturation time.</li> </ul>
<ul style="list-style-type: none"> <li>Titrate template amount. Template DNA may be damaged. Use freshly prepared template.</li> </ul>	<ul style="list-style-type: none"> <li>Check the purity and concentration of the primers.</li> <li>Check primer design.</li> </ul>
Non-specific products - High molecular weight smears	
<ul style="list-style-type: none"> <li>Make sure the extension time used was not too long.</li> </ul>	<ul style="list-style-type: none"> <li>Titrate template amount.</li> </ul>
<ul style="list-style-type: none"> <li>Reduce the total number of cycles.</li> </ul>	<ul style="list-style-type: none"> <li>Optimize denaturation temperature.</li> </ul>
<ul style="list-style-type: none"> <li>Increase annealing temperature or try 2-step PCR protocol.</li> </ul>	<ul style="list-style-type: none"> <li>Decrease primer concentration.</li> </ul>
Non-specific products - Low molecular weight discrete bands	
<ul style="list-style-type: none"> <li>Increase annealing temperature</li> </ul>	<ul style="list-style-type: none"> <li>Titrate template amount.</li> <li>Decrease primer concentration.</li> </ul>

- Make sure the extension time used was not too long.
  - Design new primers.

Component	Volume
Control DNA template	0.5 $\mu\text{l}$
Control primer Mix	0.5 $\mu\text{l}$
10mM dNTPs	0.5 $\mu\text{l}$
G9 Taq DNA Polymerase	0.25 $\mu\text{l}$
10 X Taq Reaction Buffer with $\text{MgCl}_2$	2.5 $\mu\text{l}$
Nuclease-free water	Upto 25 $\mu\text{l}$

### Control PCR cycling conditions

Cycling conditions	
Step 1	94°C – 5 min
Step 2	94°C – 30s
Step 3	57°C – 30s
Step 4	72°C – 4 min
Step 5	72°C – 10 min
Step 6	4°C – Hold

Repeat Steps 2 to 4 for 25 Cycles. A ~2kb band will be obtained in Control PCR reaction.